

Damped Lyman-Alpha Galaxies

D. Turnshek, S. Rao

Dept. of Physics and Astronomy, The University of Pittsburgh

W. Lane

Kapteyn Astronomical Institute

E. Monier

Ohio State University

D. Nestor

Dept. of Physics and Astronomy, The University of Pittsburgh

J. Bergeron

ESO Garching

A. Smette

NASA GSFC

Abstract.

Some results from an imaging program to identify low-redshift ($0.09 < z < 1.63$) damped Ly α (DLA) galaxies are presented. The standard paradigm that was widely accepted a decade ago, that DLA galaxies are the progenitors of luminous disk galaxies, is now being seriously challenged. The indisputable conclusion from imaging studies at low redshift is that the morphological types of DLA galaxies are mixed and that they span a range in luminosities and surface brightnesses.

1. Introduction

Quasar damped Ly α (DLA) absorbers are widely recognized as important probes of galaxy formation because they contain the bulk of the cosmological mass of observable neutral gas in the Universe. Consequently, studies of “DLA galaxies” are potentially among the most important tools for directly studying the conversion of gas into stars on a cosmological scale. In particular, quasar absorption-line analysis, coupled with the results from follow-up galaxy identification studies, uniquely offers the opportunity to simultaneously investigate the gaseous and stellar components of cosmologically distant galaxies, something which optical/IR galaxy surveys by themselves do not directly permit. The standard paradigm for over a decade has been that DLA systems arise in luminous HI disk galaxies, but there are now direct results which contradict this. Observations are more consistent with the idea that DLA absorption arises in Giant Hydrogen Clouds associated with various types of galaxies or protogalaxies, before the complete conversion of gas into stars (Khersonsky & Turnshek 1996).

TABLE 1

QSO	z_{DLA}	Extent ^{1,2} kpc	Luminosity ^{1,3}	Galaxy Type
0738+313	0.091	14	$M_K \approx -21.8$ (0.08 L_K^*) ⁴	LSB
	0.221	13	$M_K \approx -22.0$ (0.1 L_K^*)	dwarf
0827+243	0.518	22	$M_K \approx -24.5$ (1.0 L_K^*)	luminous spiral?
0952+179	0.239	24	$M_J \leq -19.5$ (0.03 L_J^*) ⁵	patchy/LSB?
1127-145	0.313	60	$M_J \approx -20.0$ (0.05 L_J^*) ⁵	patchy/irreg/LSB

¹For a $\Lambda = 0$ cosmology with $H_0 = 65 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $q_0 = 0.5$. ²Measured $\approx 2\sigma$ above the background. ³K-band K correction from Cowie et al. (1994, ApJ, 434, 114). $M_K^* = -24.5$ adopted from Loveday (2000, MNRAS, 312, 557), where $J - K \approx 1$ and $B - K \approx 3.5$ (Fioc & Rocca-Volmerange 1999, A&A, 351, 869). ⁴Luminosity estimated after subtraction of quasar nuclear and host light. ⁵Total luminosity estimated by summing the luminous regions not lying on the QSO.

2. Results

Recent imaging studies with *HST*-WFPC2 have begun to show that the host galaxies of confirmed low-redshift DLA absorption lines have a wide variety of morphological types. In one case, towards 3C336, there is no optical evidence for the DLA galaxy despite very deep observations (Steidel et al. 1997). In other cases (Le Brun et al. 1997) the DLA galaxies are likely to be spirals (towards 3C196 and Q1209+107), amorphous, low surface brightness galaxies (towards PKS1229-021 and 3C286), and compact objects (towards EX0302-223 and PKS0454+039). These galaxies have redshifts in the interval $0.395 \leq z \leq 1.010$.

Here we present further evidence for this mixed population from ground-based infrared observations of five of the lowest redshift DLA systems discovered in the survey of Rao & Turnshek (2000; see also Rao, these proceedings). Table 1 summarizes the properties of these five DLA galaxies and Figure 1 shows the images. Other pieces of evidence which challenge the luminous-disk hypothesis include: (1) the low-redshift DLA HI column density distribution does not fall off as $\sim N^{-3}$ at high column density, as would be expected for disks (Rao & Turnshek 2000), (2) the elemental abundances are apparently not evolving toward values typical of the solar neighborhood (Pettini et al. 1999), and (3) the kinematic properties of metal lines associated with high-redshift DLA systems can reasonably be interpreted as fragments of infalling gas (Haehnelt et al. 1998).

References

- Haehnelt, M., Steinmetz, M., & Rauch, 1998, ApJ, 495, 64
 Khersonsky, V. K., & Turnshek, D. A. 1996, ApJ, 471, 657
 Le Brun, V., Bergeron, J., Boissé, P., & Christian, C. 1993, A&A, 279, 33
 Pettini, M., Ellison, S., Steidel, C., & Bowen, D. 1999, ApJ, 510, 576
 Rao, S. M. & Turnshek, D. A. 2000, ApJS, 130, 1
 Steidel, C. C. et al. 1997, ApJ, 480, 568




Figure 1. Four QSO fields with 5 DLA systems. The light at the position of the QSOs (labeled Q) has been subtracted. North is up and east is to the left. **Top left:** IRTF K band image ($\sim 18'' \times 18''$) of the Q0738+313 field. The $z = 0.091$ DLA galaxy is identified to be the LSB galaxy nearly centered on the QSO. The dwarf galaxy $\approx 6''$ to the southeast of the QSO has a measured redshift of $z = 0.22$. See Rao & Turnshek (1998, ApJ, 500, L115) and Turnshek et al. (2000, astro-ph/0010573) for results on this field. **Top right:** IRTF K band image ($\sim 20'' \times 20''$) of the Q0827+243 field. The luminous galaxy $\approx 6''$ to the east-northeast has a measured redshift of $z = 0.52$ and is a probable spiral. Faint emission due north of the QSO could be due to another galaxy although the possibility of residuals from the QSO point spread function cannot be ruled out. **Bottom left:** NTT J band image ($\sim 24'' \times 24''$) of the Q0952+179 field. The blobs to the east and southwest of the QSO are LSB, but it is unclear if this is a patchy/irregular structure or a single LSB galaxy. **Bottom right:** NTT J band image ($\sim 27'' \times 27''$) of the Q1127-145 field. The DLA galaxy is associated with the patchy/irregular LSB (north-to-south elongated) structure $\approx 4''$ to the west; the brightest blob in this region has a measured redshift of $z = 0.31$.